


CDI POCKET GUIDE™

..... ICD-10-AM



Pinson&Tang

 UPLiftGroup

Welcome to the *CDI Pocket Guide™ ICD-10-AM*, the first Australian edition of the best-selling *CDI Pocket Guide™* that was originally created for hospitals in the United States.

That initial Guide had a long gestation period. As a consultant, Cynthia Tang witnessed first-hand the ineffectiveness of typical CDI training in the 1990's: staff would be given unwieldy three-ring binders containing hundreds of pages of information, which were difficult to search through and even more difficult to learn from. The only way to gain expert knowledge was to work in a hospital that was willing to hire consultants like her to train the staff. When Cynthia met Dr. Richard Pinson in 2006, they realised they shared a vision for helping hospitals achieve accurate and thorough documentation by explaining the principles behind the practices in such a way that everyone could easily grasp them. They decided to join forces and wrote the first *CDI Pocket Guide™* in 2007. It met with resounding success, and they have updated the Guide every year since.

As the field of CDI has emerged in Australia, Jenny Fitzpatrick and Sam Heyneman of the Uplift Group recognised the need for a resource for the Australian context. When they found the original *CDI Pocket Guide™*, they reached out to Cynthia and Richard, and together these four seasoned experts created the *CDI Pocket Guide™* for ICD-10-AM and AR-DRGs that you now hold in your hands.

This new guide retains all the clinical content that made the original *CDI Pocket Guide™* a success, but it has been revised for the ICD-10-AM classification and Australian-Refined DRGs (AR-DRGs).

We trust that you, like thousands before you, will find this a useful tool in addressing the daily complexities of coding and clinical documentation. The ultimate goal is not just more accurate coding and reimbursement, but improved quality and outcomes for both clinicians and hospitals.

To your success,
Jenny Fitzpatrick, Sam Heynemann, Richard Pinson, MD, and
Cynthia Tang

INTRODUCTION TO THE CDI POCKET GUIDE, ICD-10-AM

We trust that you will find this a useful tool in addressing the daily complexities of coding and clinical documentation.

Each of the five sections of this guide is written with a specific purpose in mind:

Guidelines is a shortcut to the most important guidelines and coding rules for DRG assignment and other important topics. Refer to these guidelines frequently.

Key References provides detailed clinical definitions and criteria, treatment, coding and documentation challenges, and references for the most important and frequently encountered conditions.

Complex Diagnoses lists and summarises the most common and important additional diagnoses with DCL that affect DRG assignment in the different AR-DRG versions.

DRG Tips includes alternative DRG assignment for select DRGs which in our experience have a high likelihood of another principal diagnosis, additional complex diagnosis or procedure.

AR-DRG Table is a complete list of the v10.0 AR-DRGs and their cost weights for quick reference.

All references and citations for The International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, Australian Modification (ICD-10-AM), Australian Classification of Health Interventions (ACHI) and Australian Coding Standards (ACS) are sourced from The Independent Hospital Pricing Authority (IHPA) 2019, Eleventh Edition publications.

References to the ACS have been shortened for simplicity to: ACS [number][Title]. References to two national coding advice sources are cited as follows: Australian Consortium for Classification Development as ACCD [date]:[Title] and Australian Classification Exchange as ACE [date]:[Title].

GUIDELINES	1	Difficult Airway/Intubation	113
Coding Rules—Overview	1	Electrolyte Disorders	116
Coding Rules—Principal Diagnosis	2	Heart Failure	119
Coding Rules—Additional Diagnosis	11	High Flow Oxygen Therapy	122
<i>Chronic Conditions (U codes)</i>	14	HIV/AIDS	124
<i>Diabetic Complications</i>	15	Intellectual Disability / Impairment	127
Coding and AR-DRG Fundamentals	18	Malnutrition	129
Condition Onset Flag (COF)	26	Microsurgical Repair	135
Ethical Queries	28	Myocardial Ischaemia/Infarction	137
Procedural Complications	36	Neoplasms	141
Quality and Safety Measures	39	Obesity and Body Mass Index	145
<i>Hospital Acquired Complications (HAC)</i>	39	Pancreatitis	148
<i>Sentinel Events</i>	43	Pathological Fractures	151
<i>Unplanned Readmissions</i>	44	Pneumonia	154
Signs, Symptoms & Unspecified Codes	46	Pressure Ulcer/Injury	159
Unrelated DRGs	49	Quadriplegia/Paraplegia	162
What Documentation Counts?	50	Respiratory Failure	164
<i>Test Results</i>	52	Sepsis	173
		Sepsis/Risk of Sepsis—Neonatal	176
KEY REFERENCES	55	Shock	179
Acute Kidney Injury	55	SIRS—Noninfectious	181
Adhesions	59	Substance Use, Abuse, Dependence	182
Anaemia	61	Transient Ischaemic Attack (TIA)	186
Antimicrobial Resistance	65	Ventilator Support	188
Asthma/Bronchitis/COPD	68		
Atelectasis	72	COMPLEX DIAGNOSES	193
Cardiac Arrest	74	Complex Diagnoses List	194
Cardiac Arrhythmia	76	Quick Reference—Interventions	203
Cerebral Oedema/Compression	80	BMI Table	204
Cerebral Palsy	82		
Cerebral Infarction/Haemorrhage	84	DRG TIPS	205
Chronic Kidney Disease	87		
Clostridium Difficile Colitis	91	AR-DRG TABLE	247
Coagulopathy Due To Anticoagulants	93		
Coma / Loss of Consciousness	96		
Cystic Fibrosis	99		
Deconditioning	101		
Delirium / Acute Confusional State	103		
Depression/Bipolar Disorder/Anxiety	105		
Diabetic Complications	108		

DEFINITION

A Transient Ischaemic Attack (TIA) is a transient episode of focal neurological dysfunction caused by cerebral ischaemia (reduced blood flow). Typical neurological deficits include focal weakness, impaired speech, difficulty with balance or walking.

DIAGNOSTIC CRITERIA

TIA is defined by both:

- Transient focal neurological deficit lasting < 24 hours, and
- No acute infarction or haemorrhage on imaging

Duration is counted from onset, not presentation. Patients often present having already had symptoms for several hours or more.

Persistence of a focal neurological deficit > 24 hours from onset is a stroke (CVA), not TIA, even with negative imaging.

While the duration of symptoms required to establish a diagnosis of TIA officially remains < 24 hours, it is a subject of debate amongst neurologists with many experts suggesting it should only be one hour.

Causes of TIA include:

- **Cerebral/pre-cerebral stenosis:** Any degree of stenosis may cause a TIA. “Noncritical stenosis” indicates that medical therapy is preferred to surgery; it does not exclude stenosis as the cause.
- **Transient cerebral embolism** due to platelet aggregates is often caused by atrial fibrillation, abnormal heart valves, or atrial septal defect (ASD).
- **Vertebrobasilar syndrome** is considered nothing more than a TIA symptom, but documentation of stenosis, occlusion, thrombosis, embolism of any vertebrobasilar arteries is a serious cause.

Common sources of transient cerebral embolism include:

- Atrial fibrillation—especially if PT/INR is subtherapeutic
- Valvular heart disease (aortic or mitral valves)

- ASD with “paradoxical embolus” of small clots from the right (venous) side of the heart across the ASD to the left heart and from there up to the brain
- Mural (ventricular wall) thrombus—especially following MI
- Any degree of stenosis/narrowing of a carotid, vertebral, or cerebral artery

Diagnostic testing often includes trans-oesophageal echocardiogram (TOE), carotid or transcranial Doppler, CT, MRI, and MRA—all looking for sources of emboli, stenosis, occlusion, or thrombosis.

TREATMENT

Treatment includes antiplatelet therapy, such as aspirin, dipyridamole or clopidogrel, to prevent recurrent thrombosis or embolism, even when no significant abnormality is identified. Less often, an anticoagulant like warfarin may be prescribed.

CODING AND DOCUMENTATION CHALLENGES

TIA is a symptom of a significant underlying cerebrovascular process. It is the underlying condition that really matters and is treated, not the symptom. Always seek clarification of the suspected or confirmed underlying cause of TIA symptoms. Even if nothing is found on evaluation to explain the TIA, the most likely cause of TIA is unexplained “transient cerebral embolism.”

Identification of CVA when criteria are actually met represents a query opportunity, especially when imaging is negative but symptoms persist > 24 hrs from onset. For **DRG Tips**, see B69 *TIA and precerebral occlusion*.

References:

- National Stroke Association Guidelines for the Management of Transient Ischemic Attacks. *Ann Neurol* 2006; 60: 301–313
- Diagnosis and Management of Transient Ischemic Attack, Coultts SB
- UpToDate.com: Initial evaluation and management of transient ischemic attack and minor ischemic stroke.

DRG H07 **Open Cholecystectomy**
(A) Major, (B) Intermediate, (C) Minor

H07A	5.9622	H02A	8.6384
H07B	3.6628	H02B	4.8246
H07C	2.5169	H02C	2.1901

PRINCIPAL DIAGNOSIS

LOOK FOR evidence of pancreatitis. Gallstone obstruction of the pancreatic or common bile duct (gallstone pancreatitis). See *Pancreatitis*.

ADDITIONAL DIAGNOSIS

LOOK FOR division of adhesions with evidence of a history of abdominal surgery to query for postprocedural adhesions. Adhesions confirmed due to previous surgery are assigned K91.89 postprocedural disorders of digestive system + K66.0 Peritoneal adhesions + appropriate external cause codes. See *Adhesions* and *Procedural Complications*.

PROCEDURE

LOOK FOR exploration of common bile duct or choledochotomy for DRG H02 (Major biliary tract interventions).

LOOK FOR evidence of adhesiolysis. Synonymous terms include 'divided', 'take down', 'released', 'dissection/dissected', 'stripped down', 'separated' or 'freed'. See *Adhesions*.

MBS item number 30393, 30378 (most frequent) are commonly documented by the surgeon without qualifying terms describing the procedure within the body of the operation report.